

Integrating trauma-informed approaches into obstetrics and gynecology medical education: a framework for safer learning and care



Celeste S. Royce, MD; Tammy Sonn, MD; Laura Baecher-Lind, MD, MPH; Katherine T. Chen, MD, MPH; Angela Fleming, DO; Christine Kim, MD; Silka Patel, MD; Hedwige Saint Louis, MD, MPH; Jonathan Schaffir, MD; Shireen Madani Sims, MD; Alyssa Stephenson-Famy, MD; Jill M. Sutton, MD; on behalf of the Undergraduate Medicine Education Committee, Association of Professors Of Gynecology And Obstetrics

Trauma-informed medical education extends the principles of trauma-informed care into medical training to foster awareness and sensitivity to trauma, aiming to avoid further harm. Trauma-informed care is patient-centered, emphasizes shared decision-making, and promotes safety, empowerment, and healing within physician–patient relationships. Similarly, trauma-informed medical education promotes a safe learning environment in which learners and educators share mutual respect and responsibilities for achieving educational goals. Trauma is a common human experience, with substantial portions of the population experiencing adverse events. Medical students also report high rates of childhood trauma. Medical education itself can exacerbate trauma, affecting learners' cognitive load, emotional resilience, and capacity to learn. The sensitive nature of obstetrics and gynecology makes it particularly relevant for a trauma-informed approach. Despite this need, trauma-informed training is sparse in medical education. Notably, few trauma-informed care curricula focus on obstetrics and gynecology, especially at the clerkship level. Incorporating trauma-informed medical education in obstetrics and gynecology curricula can enhance understanding of gynecologic and obstetric conditions' trauma-related pathophysiology. Clinical training should involve trauma-informed patient interactions, sensitive history taking, and examinations. Beyond obstetrics and gynecology, trauma-informed medical education should permeate the entire medical educational continuum, shaping academic and administrative cultures. Developing and studying the impact of trauma-informed medical education will support both learners and the patients they serve.

Key words: curriculum, medical education, obstetrics and gynecology, patient care, patient safety, trauma-informed care, trauma-informed medical education

Introduction to trauma-informed medical education

Trauma-informed medical education (TIME) is an extension of the trauma-

informed movement, which developed following the inclusion of post-traumatic stress disorder as a condition in the *Diagnostic and Statistical Manual of*

Mental Disorders (third edition; American Psychiatric Association, 1980). Trauma-informed care (TIC) espouses approaching health issues with an

From the Associate Professor and Director of Undergraduate Medical Education, Department of Obstetrics, Gynecology and Reproductive Biology, Harvard Medical School, Beth Israel Deaconess Medical Center, Boston, MA (Royce); Professor, Vice-Chair for Education, Associate Dean of Student Affairs Department of Obstetrics and Gynecology, WashU Medicine, St. Louis, MO (Sonn); Professor and Dean for Educational Affairs, Tufts School of Medicine, Boston, MA (Baecher-Lind); Professor, Obstetrics, Gynecology, and Reproductive Science and Medical Education and System Vice-Chair of Ob-Gyn Education, Icahn School of Medicine at Mount Sinai, New York, NY (Chen); Assistant Professor, Residency Program Director, Department of Obstetrics and Gynecology, Corewell Health Michigan State University College of Osteopathic Medicine, East Lansing, MI (Fleming); Health Science Clinical Professor, Assistant Residency Program Director of Transition, and Associate Clerkship Director at the Department of Obstetrics and Gynecology at the University of California, Irvine, CA (Kim); Clinical Associate Professor of Gynecology and Obstetrics at The Johns Hopkins School of Medicine, Assistant Dean of Medical Student Affairs at The Johns Hopkins School of Medicine, Baltimore, MD (Patel); Associate Professor and Clerkship Director, Department of Obstetrics & Gynecology, Morehouse School of Medicine, Atlanta, GA (Louis); Professor and Vice Chair of Education, Department of Obstetrics and Gynecology, Ohio State University Wexner Medical Center, Columbus, OH (Schaffir); Professor and Chair, Department of Obstetrics and Gynecology, University of Florida College of Medicine, Gainesville, FL (Sims); Professor Department of Obstetrics and Gynecology, University of Washington, Seattle, WA (Stephenson-Famy); and Clinical Associate Professor and Director of Undergraduate Education, Department of Obstetrics and Gynecology, Brody School of Medicine at East Carolina University Greenville, Greenville, NC (Sutton).

Received Aug. 23, 2025; revised Oct. 19, 2025; accepted Oct. 26, 2025.

The authors report no conflict of interest.

Corresponding author: Tammy Sonn, MD. sonnt@wustl.edu

0002-9378

© 2025 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

<https://doi.org/10.1016/j.ajog.2025.10.033>



Click [Supplemental Materials](#) and [Video](#) under article title in Contents at ajog.org

awareness and sensitivity to potential prior traumatic exposures, with an intention to avoid further harm.¹ TIC is an approach to patient care which is patient-centered, based on shared decision-making, and which promotes safety, empowerment, and healing within physician–patient relationships. Essential to TIC is interrupting cycles of mistrust and retraumatization, such that patients are empowered to access care on their terms, and physicians become a trusted partner in achieving health goals. Applications of TIC now extend across medical specialties. Patient satisfaction and outcomes are improved within organizations and at the individual level when TIC is practiced routinely.^{2,3} TIC espouses 6 key principles: ensuring safety for all (psychological and physical); building and maintaining trust through decisions made mutually and with transparency; peer support for healing and recovery; collaboration and mutuality with leveling of hierarchies; ensuring empowerment of all, with opportunities for expression and self-determination (“voice and choice”); and an acknowledgment and consideration of the historical, cultural, and gender-based experiences with trauma that affect individuals and communities (Table).

In the original description of TIC, an emphasis was placed on institutional support for TIC; that is, the culture of an institution needs to not only accommodate a patient-centered and trauma-informed practice but infrastructure must also be cultivated to support this practice. Part of this infrastructure development is ensuring the philosophy extends to medical education. Learners need to understand the principles of TIC as part of their training to care for patients of their own and to protect themselves against trauma experienced in the educational process itself.

A trauma-informed approach to medical education—TIME—espouses trusting relationships between learners and instructors and a safe and welcoming learning environment across the educational continuum. In turn,

multidirectional learning occurs, retraumatization is mitigated, and resiliency is promoted.⁴ In TIME, trauma-informed principles are woven throughout a curriculum: from the neuroscience and pathophysiology of trauma exposure in basic sciences, to the application of trauma-informed principles in patient care during clinical courses, and to self-care and wellness of learners to reinforce strengths and build resiliency. Applying trauma-informed principles to educating learners has the potential in turn to positively impact learners’ interactions with patients. Crucially, trauma-informed principles need to encompass the academic and administrative structure of a medical education program to fully express the benefits of TIME. In this article, we explore the literature on TIME and describe best practices to promote the incorporation of trauma-informed principles into clinical teaching in obstetrics and gynecology (OBGYN) medical education across the educational continuum.

Background of trauma and medical education

Trauma—“an event, series of events, or set of circumstances experienced by an individual as physically or emotionally harmful, or life-threatening, and that has lasting adverse effects on the individual’s functioning, and mental, physical, social, emotional, or spiritual well-being”¹—is a common human experience. Surveys consistently demonstrate rates of trauma exposure approaching 90% for adults in the United States, including a community-based survey of adults in Detroit from 1996⁵ and a nation-wide survey from 2013,⁶ while data from the World Health Organization show more than 70% of adults worldwide report lifetime experience of trauma.^{7,8} Medical students matriculate with levels of exposure to adverse childhood events (ACEs) similar to that of the general population, with 51% reporting at least one ACE and 12% reporting more than 4 ACEs,^{7,9,10} compared with general population rates of 61.6% and 15.8%,

respectively.¹¹ Although newly matriculated medical students may have greater resiliency than peers, with lower reported rates of depression, anxiety, and burnout symptoms than age-matched peers,^{9,12} the experience of medical training results in higher rates of those symptoms by the clinical years.^{13–16} While trauma exposure can and does affect people from all cohorts, spanning societal strata, race, and gender minorities, individuals from marginalized communities or low socioeconomic status have higher rates of exposure to trauma.^{6,17} Students from minoritized and marginalized backgrounds may also be at higher risk of suffering the impact of trauma¹⁸; this impact can be magnified by historical and current bias and discrimination,¹⁹ and by the hidden curriculum,²⁰ which can be described as the implicit messages, values, and community norms which are not explicitly taught as part of an educational program but which are conveyed nonetheless, and which can serve to reinforce biases and discrimination. Medical education itself can be traumatic, from mistreatment and harassment at all phases of training to vicariously experiencing traumatic events in clinical care. Learners in health professions with a prior history of trauma are at risk of retraumatization.²¹ Trauma can elevate stress levels and cognitive load, which reduces the brain’s capacity to focus and process new learning.²² Exposure to trauma may also result in learners shutting down emotionally, suffering burnout, and empathy fatigue.²³ All these can adversely impact learning capacity, learner success, and ultimately patient care.

Practicing TIC promotes health equity and access to healthcare.^{24,25} Multiple studies demonstrate the value of TIC in improved patient–clinician communications and improved engagement with healthcare.^{26,27} Retraumatization of trauma survivors may be avoided by embracing TIC.^{28,29} Application of similar principles in medical education may help ensure engagement of

TABLE
TIME in an OBGYN clerkship: examples from SAMHSA's 6 principles

Principle	Individual-level examples	Clerkship-level examples
Safety Physical and emotional safety for students and faculty; minimizing risk of retraumatization	Providing a private sleep room while on call and secure spaces for personal belongings while in clinic.	Posting face sheets of students, faculty, residents, and nurses with names, photos, and short bios on labor floor to promote team members' sense of belonging and recognition.
Trustworthiness & transparency Clear, honest communication; transparency in decision-making and procedures	Students and preceptors receive accurate schedules for clinics where students are expected to fill meaningful patient care roles.	Clerkship Director posts clear grading expectations and performance objectives online and in administrative office for learners and faculty.
Peer support Inclusion of individuals with lived experience; fostering mutual support	Student near-peer mentoring, "big sib-little sib" programs	Clerkship hosts breakfast for current students and residents for small group discussions to process clinical experiences focused on guided reflections and building resilience.
Collaboration & mutuality Shared decision-making; leveling power differences between faculty and learners	Student and preceptor together set learning goals at the start of a session or case, encouraging self-directed learning.	Students and recent graduates invited to co-create curricular content and co-teach didactics, promoting learner agency.
Empowerment, voice, & choice Supporting autonomy; validating strengths; offering choices in learning	Students choose which subspecialty clinics to attend, tailoring learning experiences to personal goals or preferences	Clerkship provides opportunities to participate in advocacy and community engagement work as option for OBGYN clerkship duties.
Recognition of cultural, historical, & gender issues Recognizing and addressing cultural, historical, and gender factors that influence trauma	Students and faculty invited to share personal stories which illuminate and influence clinical experiences, fostering identity-safe spaces.	Didactics include discussion of the role of historical and cultural disparities in disease epidemiology and adverse outcomes in OBGYN.

OBGYN, obstetrics and gynecology; SAMHSA, Substance Abuse and Mental Health Services Administration; TIME, trauma-informed medical education.

learners with educators and avoidance of retraumatization of learners in medical training.

The current state of trauma-informed medical education in obstetrics and gynecology

The relevance of a trauma-informed approach and universal application of TIC is of particular importance in the setting of OBGYN. Experiences such as sexual assault or pregnancy loss are very common throughout the population, including among patients and among medical learners. The sensitive nature of OBGYN is fraught with potential for traumatization. For medical students, the intimate nature of OBGYN histories and physical examinations can be traumatizing by reliving their own experiences. Fear of harming or hurting patients and issues of consent for examinations for educational reasons all contribute to a need for a trauma-informed approach in the learning environment. While patients welcome and expect physicians to ask about

ACEs and other trauma, physicians report discomfort with a trauma disclosure.³⁰

Although trauma exposure is ubiquitous, training in TIC is not. Gerber et al noted that of 11 published medical school curricula focused on TIME, only 2 were intended for use at the clerkship level, with 9 aimed at the preclerkship learner. A recent scoping review looking at TIME curricula for OBGYN found only one published curriculum, which was for graduate medical education learners.^{31,32}

Best practices for incorporating trauma-informed medical education into obstetrics and gynecology clerkships and advanced electives

At the basic sciences level, the role of chronic stress in the development of fibroids,³³ endometriosis,³⁴ preterm birth,³⁵ low birthweight,³⁶ and early onset preeclampsia³⁷ should be included in pathophysiology, with discussion of trauma as a contributing factor to chronic stress. This might

include an exploration of the relationship between psychological stress and oxidative stress at the cellular level and an understanding of the intergenerational effects of trauma exposure as demonstrated in epigenetic phenomena.³⁸ In clinical epidemiology and social medicine courses, instructors may examine the role of trauma exposure in chronic disease and the role of violence and abuse in women's health, for example, identifying intimate partner violence as a leading cause of maternal/fetal death.^{39,40}

In clinical clerkships, TIME can be implemented at the individual and clerkship-wide level. While single session training programs have been described,^{41,42} weaving a trauma-informed approach to patient care throughout clinical training is likely to result in more long-term familiarity and increased use of TIC. The role of TIME is perhaps best understood in the context of physical examination and clinical skills curricula. Demonstrations of TIC behaviors and language with

standardized patients, instruction in responding to trauma disclosures, and juxtaposition of trauma-informed behaviors with nontrauma-informed behaviors are useful learning tools.⁴³ Other trauma-informed approaches include explicit instruction in techniques of TIC and using a trauma-informed perspective in learner–instructor interactions. In using the TIC framework of 6 principles, Table gives specific examples of how to incorporate this on an individual and clerkship level.

Next steps in the incorporation of trauma-informed medical education into obstetrics and gynecology medical education

Resiliency

TIME provides learners with opportunities to develop resiliency and strength to respond effectively to trauma exposures, which may occur at any point in medical training.^{44–46} Trauma exposure may take many forms, including natural disasters or political events, witnessing harm occurring to a patient or population, caring for critically ill patients, or bullying. Jackson et al found as many as 57% of surgical trainees screened positive or “at risk” of post-traumatic stress disorder due to work-related experiences.⁴⁵ Application of

trauma-informed principles to medical education may mitigate the effects of this level of stress: Elsseou identified the Substance Abuse and Mental Health Services Administration’s 4-R approach (Figure) as a model for building resiliency and prevention of burnout in medical professionals, noting “TIC can transform the experience of healthcare for patients, trainees, and employees alike”.⁴⁷ Providing our learners with the skills to mitigate adverse effects of trauma is vital to help learners to cope with traumatic events of a medical education.⁴⁸ Brown et al describe “trauma stewardship”, wherein physicians manage the occupational exposure to trauma by caring for patients and understand their lived experiences without suffering the burden of that trauma themselves.³¹ In offering resources to protect learners from traumatic educational experiences, it is important not to give the impression that clinical experiences are necessarily triggering. Fostering resilience should include setting realistic expectations about the spectrum of human experience encountered, which may expand the learner’s perspective without inducing trauma.


Training faculty

Training materials aimed at practicing physicians are lacking. The lack of TIC training for faculty impacts their ability to not only embrace TIC in their own practices but also to model, teach, and assess these skills for learners. American College of Obstetricians and Gynecologists recognized in a 2021 Committee Opinion the need for trauma-informed training.⁴⁹ Brown et al in 2021 set out principles for introducing TIC into medical education, calling on medical educators to “address the epidemic of trauma and adversity faced by medical students, the medical professionals they will become, and ultimately the patients they will serve.”⁵⁰ Despite this broad need, a substantial gap exists between the recognized importance of TIC training and the implementation of structured TIC curricula. Brown et al found most TIME curricula focus on undergraduate medical education or graduate medical education, with a paucity of continuing medical education efforts, and call attention to this generational gap as amplifying the lack of TIME.³¹ While learners may be well-versed in the language and practice of TIC, practicing clinicians may not, thus contributing to the hidden curriculum and potentially to alienation and burnout of learners.

Learner participation

A key principle of TIME, based on the principles of trustworthiness, transparency, collaboration, and empowerment, is inclusion of learners in the design and implementation of curricular materials, including assessment tools. Millham et al describe a team-based approach to adapting the core Entrustable Professional Activities for Entering Residency (EPAs)⁵¹ to include trauma-informed principles, allowing faculty to give detailed assessments based on direct observation.⁴ EPAs, defined by the Association of American Medical Colleges as observable, measurable tasks performed by physicians, that trainees become entrusted to perform unsupervised once they have attained sufficient specific competence. While each EPA includes elements of Interpersonal and Communication Skills

FIGURE
The 4 Rs of trauma-informed medical education



and Professionalism, Millham et al assembled a multidisciplinary team of students, residents, and faculty to work together in an iterative, consensus-building process to modify the existing EPA framework to include assessing students' knowledge, behaviors, and skills for trauma-informed practice. Lee et al found curricular co-creation and co-implementation increases medical student engagement and satisfaction with their education.⁵² With current learners (and future educators) deeply involved in content development, TIME has the potential to improve medical school curricula to better meet the evolving needs of the profession.

Overcoming barriers

Barriers to introducing TIME may include lack of faculty with content knowledge, lack of time in an overcrowded curriculum, and unfamiliarity with existing guidelines. Faculty buy-in may be improved by departmental sponsorship of training through grand rounds presentations. Engaging with students, trainees, and staff to develop curricular content and implement curricular change can successfully augment faculty efforts and is in keeping with the principles of collaboration, mutuality, and empowerment.⁵³ Published resources for building curricular content include TIC core competencies developed by Berman et al, a trauma-informed approach to teaching history-taking⁵⁴ and the physical examination,⁴³ and assessment tools which incorporate trauma-informed principles into the entrustable professional activities framework.⁴ The National Collaborative on Trauma-Informed Health Care Education and Research offers resources, training, and a shared framework to help institutions overcome these barriers and integrate trauma-informed principles into the education of health professions.^{53,55}

Conclusion

TIME calls for the intentional integration of trauma science and principles throughout the medical curriculum, from foundational understanding of neurobiology and pathophysiology to

trauma-informed clinical care, and support for learner well-being and resilience. Trauma-informed principles applied throughout medical education have the potential to decrease burn-out, increase resiliency, and shape academic and institutional cultures to improve well-being for learners, faculty, staff, and patients. While principles of TIME are particularly relevant to OBGYN, trauma-informed principles should be developed, implemented, and studied across the educational continuum to better support both learners and the patients they serve. ■

REFERENCES

1. Substance Abuse and Mental Health Services. SAMHSA's concept of trauma and guidance for a trauma-informed approach. 2014. https://ncsacw.samhsa.gov/userfiles/files/SAMHSA_Trauma.pdf. Accessed February 17, 2025.
2. Kuehn BM. Trauma-informed care may ease patient fear, clinician burnout. *JAMA* 2020;323:595–7. <https://doi.org/10.1001/jama.2020.0052>.
3. Owens L, Terrell S, Low LK, et al. Universal precautions: the case for consistently trauma-informed reproductive healthcare. *Am J Obstet Gynecol* 2022;226:671–7. <https://doi.org/10.1016/j.ajog.2021.08.012>.
4. Millham LRI, Potter J, Hirsh DA, et al. Incorporation of trauma-informed care into entrustable professional activities for medical student assessment. *Acad Med* 2024. <https://doi.org/10.1097/ACM.0000000000005824>.
5. Breslau N, Kessler RC, Chilcoat HD, Schultz LR, Davis GC, Andreski P. Trauma and posttraumatic stress disorder in the community: the 1996 Detroit area survey of trauma. *Arch Gen Psychiatry* 1998;55:626–32. <https://doi.org/10.1001/archpsyc.55.7.626>.
6. Kilpatrick DG, Resnick HS, Milanak ME, Miller MW, Keyes KM, Friedman MJ. National estimates of exposure to traumatic events and PTSD prevalence using DSM-IV and DSM-5 criteria. *J Trauma Stress* 2013;26:537–47. <https://doi.org/10.1002/jts.21848>.
7. Felitti VJ, Anda RF, Nordenberg D, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The adverse childhood experiences (ACE) study. *Am J Prev Med* 1998;14:245–58.
8. Liu H, Petukhova MV, Sampson NA, et al. Association of DSM-IV posttraumatic stress disorder with traumatic experience type and history in the World health organization World mental health surveys. *JAMA Psychiatry* 2017;74:270–81. <https://doi.org/10.1001/jamapsychiatry.2016.3783>.
9. Sciolla AF, Wilkes MS, Griffin EJ. Adverse childhood experiences in medical students: implications for wellness. *Acad Psychiatry* 2019;43:369–74. <https://doi.org/10.1007/s40596-019-01047-5>.
10. Yang P, Robles-Ramamurthy B, Plastino KA. Associations between adverse childhood experiences and medical students' interest in careers: a single-setting study. *Front Psychiatry* 2025;16:1483871. <https://doi.org/10.3389/fpsy.2025.1483871>.
11. Merrick MT, Ford DC, Ports KA, Guinn AS. Prevalence of adverse childhood experiences from the 2011–2014 behavioral risk factor surveillance system in 23 states. *JAMA Pediatr* 2018;172:1038–44. <https://doi.org/10.1001/jamapediatrics.2018.2537>.
12. Blickenstaff HR, Bastin TJ, Byram JN. Exploring resilience factors in medical students with adverse childhood experiences: a pilot study. *Acad Psychiatry J Am Assoc Dir Psychiatr Resid Train Assoc Acad Psychiatry* 2022;46:218–22. <https://doi.org/10.1007/s40596-021-01560-6>.
13. Rotenstein LS, Ramos MA, Torre M, et al. Prevalence of depression, depressive symptoms, and suicidal ideation among medical students: a systematic review and meta-analysis. *JAMA* 2016;316:2214. <https://doi.org/10.1001/jama.2016.17324>.
14. Quek TTC, Tam WWS, Tran BX, et al. The global prevalence of anxiety among medical students: a meta-analysis. *Int J Environ Res Public Health* 2019;16:2735. <https://doi.org/10.3390/ijerph16152735>.
15. Dyrbye LN, West CP, Satele D, et al. Burnout among U.S. medical students, residents, and early career physicians relative to the general U.S. population. *Acad Med J Assoc Am Med Coll* 2014;89:443–51. <https://doi.org/10.1097/ACM.000000000000134>.
16. Brazeau CMLR, Shanafelt T, Durning SJ, et al. Distress among matriculating medical students relative to the general population. *Acad Med J Assoc Am Med Coll* 2014;89:1520–5. <https://doi.org/10.1097/ACM.0000000000000482>.
17. Giano Z, Wheeler DL, Hubach RD. The frequencies and disparities of adverse childhood experiences in the U.S. *BMC Public Health* 2020;20:1327. <https://doi.org/10.1186/s12889-020-09411-z>.
18. McClinton A, Laurencin CT. Just in TIME: trauma-informed medical education. *J Racial Ethn Health Disparities* 2020;7:1046–52. <https://doi.org/10.1007/s40615-020-00881-w>.
19. Matheson K, Foster MD, Bombay A, McQuaid RJ, Anisman H. Traumatic experiences, perceived discrimination, and psychological distress among members of various socially marginalized groups. *Front Psychol* 2019;10:416. <https://doi.org/10.3389/fpsyg.2019.00416>.
20. Hafferty FW. Beyond curriculum reform: confronting medicine's hidden curriculum. *Acad Med J Assoc Am Med Coll* 1998;73:

- 403–7. <https://doi.org/10.1097/00001888-199804000-00013>.
21. Li Y, Cannon LM, Coolidge EM, Darling-Fisher CS, Pardee M, Kuzma EK. Current state of trauma-informed education in the health sciences: lessons for nursing. *J Nurs Educ* 2019;58:93–101. <https://doi.org/10.3928/01484834-20190122-06>.
22. Iskander M. Burnout, cognitive overload, and metacognition in medicine. *Med Sci Educ* 2019;29:325–8. <https://doi.org/10.1007/s40670-018-00654-5>.
23. Stout J, Martin AI. Trauma-informed care in the classroom: our experience with a content warning in a medical school course. *Med Sci Educ* 2022;32:711–8. <https://doi.org/10.1007/s40670-022-01559-0>.
24. Levy-Carrick NC, Lewis-O'Connor A, Rittenberg E, Manosvalas K, Stoklosa HM, Silbersweig DA. Promoting health equity through trauma-informed care: critical role for physicians in policy and program development. *Fam Community Health* 2019;42:104–8. <https://doi.org/10.1097/FCH.0000000000000214>.
25. Tebes JK, Champine RB, Matlin SL, Strambler MJ. Population health and trauma-informed practice: implications for programs, systems, and policies. *Am J Community Psychol* 2019;64:494–508. <https://doi.org/10.1002/ajcp.12382>.
26. Chaudhri S, Zweig KC, Hebbar P, Angell S, Vasan A. Trauma-informed care: a strategy to improve primary healthcare engagement for persons with criminal justice system involvement. *J Gen Intern Med* 2019;34:1048–52. <https://doi.org/10.1007/s11606-018-4783-1>.
27. Osofsky HJ, Osofsky JD, Hansel TC, Flynn T. The Louisiana mental and behavioral health capacity project trauma-informed integrated care model and improved posttraumatic stress outcomes. *J Public Health Manag Pract JPHMP* 2017;23(Suppl 6):S25–31. <https://doi.org/10.1097/PHH.0000000000000652>.
28. O'Dwyer C, Tarzia L, Fernbacher S, Hegarty K. Health professionals' experiences of providing care for women survivors of sexual violence in psychiatric inpatient units. *BMC Health Serv Res* 2019;19:839. <https://doi.org/10.1186/s12913-019-4683-z>.
29. Sweeney A, Perôt C, Callard F, Adenden V, Mantovani N, Goldsmith L. Out of the silence: towards grassroots and trauma-informed support for people who have experienced sexual violence and abuse. *Epidemiol Psychiatr Sci* 2019;28:598–602. <https://doi.org/10.1017/S2045796019000131>.
30. Goldstein E, Athale N, Sciolla AF, Catz SL. Patient preferences for discussing childhood trauma in primary care. *Perm J* 2017;21:16–055. <https://doi.org/10.7812/TPP/16-055>.
31. Burns CJ, Borah L, Terrell SM, James LTN, Erkkinen E, Owens L. Trauma-informed care curricula for the health professions: a scoping review of best practices for design, implementation, and evaluation. *Acad Med* 2023;98:401–9. <https://doi.org/10.1097/ACM.0000000000005046>.
32. Stevens NR, Holmgren L, Hobfoll SE, Cvendros JA. Assessing trauma history in pregnant patients: a didactic module and role-play for obstetrics and gynecology residents. *MedEdPORTAL J Teach Learn Resour* 2020;16:10925. https://doi.org/10.15766/mep_2374-8265.10925.
33. Qin H, Lin Z, Vásquez E, Xu L. The association between chronic psychological stress and uterine fibroids risk: a meta-analysis of observational studies. *Stress Health* 2019;35:585–94. <https://doi.org/10.1002/smi.2895>.
34. Reis FM, Coutinho LM, Vannuccini S, Luisi S, Petraglia F. Is stress a cause or a consequence of endometriosis? *Reprod Sci* 2020;27:39–45. <https://doi.org/10.1007/s43032-019-00053-0>.
35. Roy-Matton N, Moutquin JM, Brown C, Carrier N, Bell L. The impact of perceived maternal stress and other psychosocial risk factors on pregnancy complications. *J Obstet Gynaecol Can* 2011;33:344–52. [https://doi.org/10.1016/S1701-2163\(16\)34852-6](https://doi.org/10.1016/S1701-2163(16)34852-6).
36. Smith MV, Gotman N, Yonkers KA. Early childhood adversity and pregnancy outcomes. *Matern Child Health J* 2016;20:790–8. <https://doi.org/10.1007/s10995-015-1909-5>.
37. van Esch JJA, Bolte AC, Spaanderman MEA, Vandenbussche FPHA, de Weerth C, Beijers R. Maternal anxiety forecasts shorter prolongation of pregnancies complicated by early-onset preeclampsia. *Arch Gynecol Obstet* 2023;308:1703–11. <https://doi.org/10.1007/s00404-022-06836-2>.
38. Yehuda R, Lehrner A. Intergenerational transmission of trauma effects: putative role of epigenetic mechanisms. *World Psychiatry Off J World Psychiatr Assoc WPA* 2018;17:243–57. <https://doi.org/10.1002/wps.20568>.
39. Alhusein JL, Bullock L, Sharps P, Schminkey D, Comstock E, Campbell J. Intimate partner violence during pregnancy and adverse neonatal outcomes in low-income women. *J Womens Health* 2014;23:920–6. <https://doi.org/10.1089/jwh.2014.4862>.
40. Chamberlain L, Levenson R, Lee NC, et al. Addressing intimate partner violence, reproductive and sexual coercion: a seminar-style course for senior medical students. *Med Sci Educ* 2023;33:1481–6. <https://doi.org/10.1007/s40670-023-01918-5>.
41. Goldstein E, Murray-García J, Sciolla AF, Topitzes J. Medical students' perspectives on trauma-informed care training. *Perm J* 2018;22:17–126. <https://doi.org/10.7812/TPP/17-126>.
42. Gore DJ, Prusky M, Solomon CJ, et al. Creation of a medical student training to improve comfort providing trauma-informed care to sexual assault survivors. *MedEdPORTAL J Teach Learn Resour* 2021;17:11140. https://doi.org/10.15766/mep_2374-8265.11140.
43. Elisseou S, Puranam S, Nandi M. A novel, trauma-informed physical examination curriculum for first-year medical students. *MedEdPORTAL* 2019;15:10799. https://doi.org/10.15766/mep_2374-8265.10799.
44. Vance MC, Mash HBH, Ursano RJ, et al. Exposure to workplace trauma and post-traumatic stress disorder among intern physicians. *JAMA Netw Open* 2021;4:e2112837. <https://doi.org/10.1001/jamanetworkopen.2021.12837>.
45. Jackson T, Provencio A, Bentley-Kumar K, et al. PTSD and surgical residents: everybody hurts... sometimes. *Am J Surg* 2017;214:1118–24. <https://doi.org/10.1016/j.amsurg.2017.08.037>.
46. Joseph B, Pandit V, Hadeed G, et al. Unveiling posttraumatic stress disorder in trauma surgeons: a national survey. *J Trauma Acute Care Surg* 2014;77:148–54. <https://doi.org/10.1097/TA.0000000000000271>.
47. Elisseou S. Trauma-informed care: a missing link in addressing burnout. *J Healthc Leadersh* 2023;15:169–73. <https://doi.org/10.2147/JHL.S389271>.
48. Boyd JE, Lanius RA, McKinnon MC. Mindfulness-based treatments for post-traumatic stress disorder: a review of the treatment literature and neurobiological evidence. *J Psychiatry Neurosci JPN* 2018;43:7–25. <https://doi.org/10.1503/jpn.170021>.
49. Caring for patients who have experienced trauma. <https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2021/04/caring-for-patients-who-have-experienced-trauma>. Accessed February 13, 2025.
50. Brown T, Berman S, McDaniel K, et al. Trauma-informed medical education (TIME). *Acad Med* 2020. <https://doi.org/10.1097/acm.0000000000003587>.
51. Association of American Medical Colleges. Core entrustable professional activities for entering residency. *Acad Med* 2013;90:431–6. <https://doi.org/10.1097/ACM.0000000000000586>.
52. Lee AC, Gooding HC, Stoddard HA. A case study of curriculum Co-creation during a seminar-style course for senior medical students. *Med Sci Educ* 2023;33:1481–6. <https://doi.org/10.1007/s40670-023-01918-5>.
53. Berman S, Brown T, Mizelle C, et al. Roadmap for trauma-informed medical education: introducing an essential competency set. *Acad Med* 2023;98:882. <https://doi.org/10.1097/ACM.0000000000005196>.
54. Machtinger EL, Davis KB, Kimberg LS, et al. From treatment to healing: inquiry and response to recent and past trauma in adult health care. *Womens Health Issues Off Publ Jacobs Inst Womens Health* 2019;29:97–102. <https://doi.org/10.1016/j.whi.2018.11.003>.
55. TIC4MedEd.org. TIC4MedEd.org. <https://tic4meded.org>. Accessed June 29, 2025.